

# Preparing for Birth: Mothers-References

Aasheim, V. *et al.* (2017) 'Perineal techniques during the second stage of labour for reducing perineal trauma', *Cochrane Database of Systematic Reviews*, 2017(6). doi: 10.1002/14651858.CD006672.pub3.

Ahmad, A. *et al.* (2014) 'Association between fetal position at onset of labor and mode of delivery: A prospective cohort study', *Ultrasound in Obstetrics and Gynecology*, 43(2), pp. 176–182. doi: 10.1002/uog.13189.

Alfirevic, Z. *et al.* (2015a) 'Labour induction with prostaglandins: A systematic review and network meta-analysis', *BMJ (Online)*, 350. doi: 10.1136/bmj.h217.

Alfirevic, Z. *et al.* (2015b) 'Labour induction with prostaglandins: A systematic review and network meta-analysis', *Obstetrical and Gynecological Survey*, 70(6), pp. 371–373. doi: 10.1097/OGX.0000000000000213.

American College of Obstetricians and Gynecologists. (2020). Ultrasound Exams. ONLINE <https://www.acog.org/womens-health/faqs/ultrasound-exams>

Anim-Somuah, M. *et al.* (2018) 'Epidural versus non-epidural or no analgesia for pain management in labour', *Cochrane Database of Systematic Reviews*. doi: 10.1002/14651858.CD000331.pub4.

Aquino, C. I. *et al.* (2020) 'Perineal massage during labor: a systematic review and meta-analysis of randomized controlled trials', *Journal of Maternal-Fetal and Neonatal Medicine*, 33(6), pp. 1051–1063. doi: 10.1080/14767058.2018.1512574.

Association of Chartered Physiotherapists in Women's Health (2004) 'Fit and Safe to exercise in the childbearing year'. London: Author, pp. 1–20.

Augustinsson, L. E., Bohlin, P; Bundsen, P., Carlsson, C. A., Forssman, L., Sjöberg, P., Tyreman, N. O. (1977). Pain relief during delivery by transcutaneous electrical nerve stimulation. *Pain*, 4(1), 59-65.

Australian Commission on Safety and Quality in Health Care (2021a) 'Third and Fourth Degree Perineal Tears'. Sydney: Australian Commission on Safety and Quality in Health Care, pp. 1–5. doi: 10.1007/978-1-84628-503-5\_4.

Australian Commission on Safety and Quality in Health Care (2021b) *Third and Fourth Degree Perineal Tears Clinical Care Standard, Clinical Standards of Care*. Sydney. doi: 10.1136/bmj.e774.

Báez-Suárez, A., Martín-Castillo, E., García-Andújar, J., García-Hernández, J. Á., Quintana-Montesdeoca, M. P., & Loro-Ferrer, J. F. (2018). Evaluation of different doses of

transcutaneous nerve stimulation for pain relief during labour: a randomized controlled trial. *Trials*, 19(1), 652. doi:10.1186/s13063-018-3036-2

Beckmann MM, Stock OM. (2013). Antenatal perineal massage for reducing perineal trauma. *Cochrane Database of Systematic Reviews*, Issue 4. Art. No.: CD005123. DOI: 10.1002/14651858.CD005123.pub3. Accessed 11 April 2021.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD005123.pub3/full?highlightAbstract=perineal%7Cperin%7Cmassage%7Cmassag>

Beckmann, M. et al. (2017) 'Measuring women's experience of induction of labor using prostaglandin vaginal gel', *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 210, pp. 189–195. doi: 10.1016/j.ejogrb.2016.12.032.

Beckmann, M. et al. (2020) 'Induction of labour using prostaglandin E2 as an inpatient versus balloon catheter as an outpatient: a multicentre randomised controlled trial', *BJOG: An International Journal of Obstetrics and Gynaecology*, 127(5), pp. 571–579. doi: 10.1111/1471-0528.16030.

Begley CM, Gyte GM, Devane D, McGuire W, Weeks A. (2011). Active versus expectant management for women in the third stage of labour. *Cochrane Database Syst Rev*(11):CD007412. doi: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4026059/>

Begley, C. M. et al. (2012) 'Irish and New Zealand midwives' expertise in expectant management of the third stage of labour: The "MEET" study', *Midwifery*, 28(6), pp. 733–739. doi: 10.1016/j.midw.2011.08.008.

Begley, C. M. et al. (2019) 'Active versus expectant management for women in the third stage of labour', *Cochrane Database of Systematic Reviews*, 11(CD007412), pp. 1–161. doi: 10.1002/14651858.CD007412.pub5.

Berens, P., Labbok, M. and The Academy of Breastfeeding Medicine (2015) 'ABM clinical protocol #13: Contraception during breastfeeding, revised 2015', *Breastfeeding Medicine*, 10(1), pp. 3–12. doi: 10.1089/bfm.2015.9999.

Bergman, N. J. (2013) 'Neonatal stomach volume and physiology suggest feeding at 1-h intervals', *Acta Paediatrica, International Journal of Paediatrics*, 102(8), pp. 773–777. doi: 10.1111/apa.12291.

Boulvain, M., Kelly, A. J. and Irion, O. (2008) 'Intracervical prostaglandins for induction of labour', *Cochrane Database of Systematic Reviews*, (1). doi: 10.1002/14651858.CD006971. Bricker, L., Medley, N. and Pratt, J. J. (2015) 'Routine ultrasound in late pregnancy (after 24 weeks' gestation)', *Cochrane Database of Systematic Reviews*. doi: 10.1002/14651858.CD001451.pub4.

Brand, A., Hammon, I., Mountford, J., Roeske, L. & Cancer Council Australia Cervical Cancer Screening Guidelines Working Party. (2019). Screening in pregnancy. Online. [https://wiki.cancer.org.au/australia/Clinical\\_question:Screening\\_in\\_pregnancy](https://wiki.cancer.org.au/australia/Clinical_question:Screening_in_pregnancy)

Brennan, M., Clarke, M., & Devane, D. (2016). The use of anti-stretch marks' products by women in pregnancy: A descriptive, cross-sectional survey. *BMC Pregnancy and Childbirth* 16:276

Bricker, L., Medley, N., Pratt, J.J. Routine ultrasound in late pregnancy (after 24 weeks' gestation). Cochrane Database of Systematic Reviews 2015, Issue 6. Art. No.: CD001451. DOI: 10.1002/14651858.CD001451.pub4. Accessed 21 March 2021.

Campbell, S. (2013) 'A short history of sonography in obstetrics and gynaecology.', *Facts, views & vision in ObGyn*, 5(3), pp. 213–229. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24753947> <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3987368/>.

Carvalho, O. M. C. et al. (2019) 'Late umbilical cord clamping does not increase rates of jaundice and the need for phototherapy in pregnancies at normal risk', *Journal of Maternal-Fetal and Neonatal Medicine*, 32(22), pp. 3824–3829. doi: 10.1080/14767058.2018.1473367.

Cohen, S. R. and Thomas, C. R. (2015) 'Rebozo Technique for Fetal Malposition in Labor', *Journal of Midwifery and Women's Health*, 60(4), pp. 445–451. doi: 10.1111/jmwh.12352. Committee on Obstetric Practice (2017) *Methods for Estimating the Due Date*. Available at: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2017/05/methods-for-estimating-the-due-date>.

Davenport, M. H. et al. (2020) 'Infographic. Prenatal physical activity: Baby steps for better health', *British Journal of Sports Medicine*, 54(6), pp. 360–361. doi: 10.1136/bjsports-2019-101056.

Davey, M. A. and King, J. (2016) 'Caesarean section following induction of labour in uncomplicated first births- a population-based cross-sectional analysis of 42,950 births', *BMC Pregnancy and Childbirth*, 16(1), pp. 1–9. doi: 10.1186/s12884-016-0869-0.

Day, M. L. and Snell, B. J. (1993) 'Use of prostaglandins for induction of labor', *Journal of Nurse-Midwifery*, 38(2 SUPPL.), pp. 42s-48s. doi: 10.1016/0091-2182(93)90096-Y.

Dekker, R. (2018). Breathing for pain relief during labor. Accessed online: <https://evidencebasedbirth.com/breathing-for-pain-relief-during-labor/>

Department of Health (2020) Clinical Practice Guidelines: Pregnancy Care. Canberra: Australian Government Department of Health.

Dicke, J. M., Piper, S. L. and Goldfarb, C. A. (2015) 'The utility of ultrasound for the

detection of fetal limb abnormalities - a 20-year single-center experience', *Prenatal Diagnosis*, 35(4), pp. 348–353. doi: 10.1002/pd.4546.

Diederer, M. et al. (2018) 'Safety of the balloon catheter for cervical ripening in outpatient care: complications during the period from insertion to expulsion of a balloon catheter in the process of labour induction: a systematic review', *BJOG: An International Journal of Obstetrics and Gynaecology*, 125(9), pp. 1086–1095. doi: 10.1111/1471-0528.15047.

Dodd JM, Crowther CA, Huertas E, Guise JM, Horey D. (2013). Planned elective repeat caesarean section versus planned vaginal birth for women with a previous caesarean birth. Cochrane Database of Systematic Reviews, Issue 12. Art. No.: CD004224. DOI: 10.1002/14651858.CD004224.pub3.

Domenjoz, I., Kayser, B. and Boulvain, M. (2014) 'Effect of physical activity during pregnancy on mode of delivery', *American Journal of Obstetrics and Gynecology*, 211(4), pp. 401.e1-401.e11. doi: 10.1016/j.ajog.2014.03.030.

Dowswell, T., Bedwell, C., Lavender, T., Neilson, J. P. (2009). Transcutaneous electrical nerve stimulation (TENS) for pain management in labour. Cochrane Database Systematic Review, April 15(2). doi: <https://doi.org/10.1002/14651858.CD007214.pub2>

Duncan, L. G. et al. (2017) 'Benefits of preparing for childbirth with mindfulness training: A randomized controlled trial with active comparison', *BMC Pregnancy and Childbirth*, 17(1), pp. 1–11. doi: 10.1186/s12884-017-1319-3.

Evans, S. J. and Cremering, M. M. (2016) 'Use of Peanut Labor Ball for Pelvic Positioning for Nulliparous Women Following Epidural Anesthesia', *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 45(3), p. S47. doi: 10.1016/j.jogn.2016.03.119.

Fealy, S. M. et al. (2017) 'Weighing as a stand-alone intervention does not reduce excessive gestational weight gain compared to routine antenatal care: A systematic review and meta-analysis of randomised controlled trials', *BMC Pregnancy and Childbirth*, 17(1), pp. 1–12. doi: 10.1186/s12884-016-1207-2.

French, C. A., Cong, X. and Chung, K. S. (2016) 'Labor Epidural Analgesia and Breastfeeding: A Systematic Review', *Journal of Human Lactation*, 32(3), pp. 507–520. doi: 10.1177/0890334415623779.

Gommers, J. S. M. et al. (2017) 'Risk of maternal, fetal and neonatal complications associated with the use of the transcervical balloon catheter in induction of labour: A systematic review', *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 218, pp. 73–84. doi: 10.1016/j.ejogrb.2017.09.014.

Guasch, E. et al. (2017) 'Failed epidural for labor: What now?', *Minerva Anestesiologica*, 83(11), pp. 1207–1213. doi: 10.23736/S0375-9393.17.12082-1.

Guittier, MJ., Othenin-Girard, V., Irion, O. *et al.* Maternal positioning to correct occipito-posterior fetal position in labour: a randomised controlled trial. *BMC Pregnancy Childbirth* **14**, 83 (2014). <https://doi.org/10.1186/1471-2393-14-83>

Hurst, N. M. (2007) 'Recognizing and treating delayed or failed lactogenesis II.', *Journal of Midwifery & Women's Health*, 52(6), pp. 588–594. doi: 10.1016/j.jmwh.2007.05.005.

Hutton, E. K. and Mozurkewich, E. L. (2001) 'Extra-amniotic prostaglandin for induction of labour', *Cochrane Database of Systematic Reviews*. doi: 10.1002/14651858.cd003092.

Jordon R. G., Farle, C. L. & Grace, K. T. *Prenatal and Postnatal Care: A Woman-Centred Approach*, 2<sup>nd</sup> ed. (2019). John Wiley and Sons.

Jozwiak, M. *et al.* (2011) 'Foley catheter versus vaginal prostaglandin E2 gel for induction of labour at term (PROBAAT trial): An open-label, randomised controlled trial', *The Lancet*, 378(9809), pp. 2095–2103. doi: 10.1016/S0140-6736(11)61484-0.

Karim, J. N. *et al.* (2017) 'Systematic review of first-trimester ultrasound screening for detection of fetal structural anomalies and factors that affect screening performance', *Ultrasound in Obstetrics and Gynecology*, 50(4), pp. 429–441. doi: 10.1002/uog.17246.

Kayman-Kose, S., Arioz, D. T., Toktas, H., Koken, G., Kanat-Pektaş, M., Kose, M., & Yilmazer, M. (2014). Transcutaneous electrical nerve stimulation (TENS) for pain control after vaginal delivery and cesarean section. *The Journal of Maternal-Fetal & Neonatal Medicine*, 27(15), 1572-1575. doi:10.3109/14767058.2013.870549

Kinser, P. A. *et al.* (2017) 'Physical Activity and Yoga-Based Approaches for Pregnancy-Related Low Back and Pelvic Pain', *JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 46(3), pp. 334–346. doi: 10.1016/j.jogn.2016.12.006.

Kjeldsen, L. L., Sindberg, M. and Maimburg, R. D. (2015) 'Earlier induction of labour in post term pregnancies-A historical cohort study', *Midwifery*, 31(5), pp. 526–531. doi: 10.1016/j.midw.2015.02.003.

Lavender T, Hofmeyr GJ, Neilson JP, Kingdon C, Gyte GML. (2012). Caesarean section for non-medical reasons at term. *Cochrane Database of Systematic Reviews*, Issue 3. Art. No.: CD004660. DOI: 10.1002/14651858.CD004660.pub3. Accessed 23 April 2021.

Logan, L. and Stettler, S. (2016) 'Reduction of Primary Cesarean Birth Rate in a Rural Hospital Driven By Nurse-Initiated Peanut Ball Use in Active Labor', *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 48(3), p. S110. doi: 10.1016/j.jogn.2019.04.185.

McCarthy, F. P. and Kenny, L. C. (2011) 'Induction of labour', *Obstetrics, Gynaecology and Reproductive Medicine*, 21(1), pp. 1–6. doi: 10.1016/j.ogrm.2010.09.008.

Mid Essex Hospital Services NHS Trust (2016) 'Exercise and Advice Following a

Caesarean Birth', pp. 1–8. Available at:

<https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwibzLjByNLvAhVA73MBHaPIBcg4ChAWMAI6BAgJEA&url=https%3A%2F%2Fwww.meht.nhs.uk%2FEasysiteWeb%2Fgetresource.axd%3FAssetID%3D2433%26type%3Dfull%26service%3DAttachment&usg=AOvVaw1labA>.

Meisel P, Meisel M. Beeinflussen Arzneimittel in der Schwangerschaft den Neugeborenenikterus? [Do drugs in pregnancy modify neonatal jaundice?]. Zentralbl Gynakol. 1993;115(7):301-8. German. PMID: 8368029.

Mohammad, P. A. S., Doosti, C. R. and Mirghafourvand, M. (2021) 'Use of the peanut ball during labour: A systematic review and analysis', *Nursing Open*, (August 2020), pp. 1–9. doi: 10.1002/nop2.844.

Mottola, M. F. et al. (2018) '2019 Canadian Guideline for Physical Activity throughout Pregnancy', *Journal of Obstetrics and Gynaecology Canada*, 40(11), pp. 1528–1537. doi: 10.1016/j.jogc.2018.07.001.

Movementmatters.org.au (2021) 'Your Baby's Movements Matter'. Available at: <http://movementsmatter.org.au/information-for-women/>.

Mozurkewich, E. et al. (2009) 'Indications for induction of labour: A best-evidence review', *BJOG: An International Journal of Obstetrics and Gynaecology*, 116(5), pp. 626–636. doi: 10.1111/j.1471-0528.2008.02065.x.

Munim, S., Nadeem, S. and Ali Khuwaja, N. (2006) 'The accuracy of ultrasound in the diagnosis of congenital abnormalities', *Journal of the Pakistan Medical Association*, 56(1), pp. 16–18.

Nelson, T. R. et al. (2009) 'Ultrasound biosafety considerations for the practicing sonographer and sonologist', *Journal of Ultrasound in Medicine*, 28(2), pp. 139–150. doi: 10.7863/jum.2009.28.2.139.

Nichols, L. (2018). *Real Food for Pregnancy: The Science and Wisdom of Optimal Prenatal Nutrition*. USA.

Pairman, S., Pincombe, J., Thorogood, C. & Tracy, S. (2015). *Midwifery preparation for practice* 3 Ed. Sydney: Churchill Livingstone.

Pelvic Obstetric and Gynaecological Physiotherapy (2018) 'Exercise and advice after pregnancy', pp. 1–10. Available at: [https://thepogp.co.uk/\\_userfiles/pages/files/POGP-EAA5-UL.pdf](https://thepogp.co.uk/_userfiles/pages/files/POGP-EAA5-UL.pdf).

Phang-Davey, G. (2021). OMP Reversing Protocols. Accessed online <https://www.optimalmaternalpositioning.com/omp-reversing-protocols>

Polis, R. L., Gussman, D. and Kuo, Y. H. (2015) 'Yoga in pregnancy: An examination of maternal and fetal responses to 26 yoga postures', *Obstetrics and Gynecology*, 126(6), pp. 1237-1241. doi: 10.1097/AOG.0000000000001137.

Potter, K. and Brown, S. E. (2020) 'Increasing Use of the Peanut Ball in Labor Introduction', *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 49(6), p. S46. doi: 10.1016/j.jogn.2020.09.079.

Queensland Health. (2017). Opioids in labour. Queensland Clinical Guidelines. Accessed: [https://www.health.qld.gov.au/\\_data/assets/pdf\\_file/0035/682838/sg-opioids.pdf](https://www.health.qld.gov.au/_data/assets/pdf_file/0035/682838/sg-opioids.pdf)

Queensland Health (2018) 'Short Guide: Fetal movements', *Queensland Clinical Guidelines*, (October), pp. 1-5.

Raudenska, Jaroslave, and Alena Javurkova. (2016) *The Psychological Context of Labour Pain*, Nova Science Publishers, Incorporated, ProQuest Ebook Central, <https://ebookcentral.proquest.com/lib/uwsau/detail.action?docID=4586588>.

Reed, R., (2013). *Feel the fear and birth anyway*. Accessed online <https://midwifethinking.com/2013/03/27/feel-the-fear-and-birth-anyway/>

Reed, R., (2018). *Why induction matters*. London: Printer & Martin.

Roth, C. et al. (2016) 'Randomized controlled trial of Use of the Peanut Ball During Labour', *MCN, The American Journal of Maternal Child Nursing* 41(3), pp. 140-146.

Rottenstreich, M. et al. (2019) 'A rare presentation of uterine rupture—the risk of sequential labour induction with prostaglandins and oxytocin', *Journal of Obstetrics and Gynaecology*, 39(5), pp. 714-715. doi: 10.1080/01443615.2018.1553940.

Royal College of Obstetricians and Gynaecologists. (2010). Amniocentesis and Chorionic Villus Sampling. Green-Top Guideline No. 8

Santana, L. S., Gallo, R. B. S., Ferreira, C. H. J., Duarte, G., Quintana, S. M., Marcolin, A. C. (2016). Transcutaneous electrical nerve stimulation (TENS) reduces pain and postpones the need for pharmacological analgesia during labour: a randomised trial. *Journal of Physiotherapy*, 62, 29-34. doi: <https://doi.org/10.1016/j.jphys.2015.11.002>

Seijmonsbergen-Schermers, A. E., Scherjon, S. and de Jonge, A. (2019) 'Induction of labour should be offered to all women at term: AGAINST: Induction of labour should not be offered to all women at term: first, do no harm', *BJOG: An International Journal of Obstetrics and Gynaecology*, 126(13), p. 1599. doi: 10.1111/1471-0528.15887.

Seitz, J. (2017). The importance of skin-to-skin with baby after delivery. Accessed online: <https://news.sanfordhealth.org/childrens/the-importance-of-skin-to-skin-after-delivery-you-should-know/>

Shahoei, R., Shahghebi, S., Rezaei, M., Naqshbandi, S. (2017). The effect of transcutaneous electrical nerve stimulation on the severity of labor pain among nulliparous women: A clinical trial. *Complementary Therapies in Clinical Practice*, 28, 176-180. doi: <https://doi.org/10.1016/j.ctcp.2017.05.004>

Shoorab, N. J. et al. (2019) 'Women's experiences of emotional recovery from childbirth-related perineal trauma: A qualitative content analysis', *International Journal of Community Based Nursing and Midwifery*, 7(3), pp. 181–191. doi: 10.30476/IJCBNM.2019.44993.

Simmons, S. et al. (2007) 'Combined spinal-epidural versus epidural analgesia in labour', *Cochrane Database of Systematic Reviews*. doi: 10.1002/14651858.CD003401.pub2.

Souza, C. De et al. (2016) 'Length of perineal pain relief after ice pack application : A quasi-experimental study', *Women and Birth*, 29(2), pp. 117–122. doi: 10.1016/j.wombi.2015.09.002.

Smith CA, Levett KM, Collins CT, Armour M, Dahlen HG, Suganuma M. (2018). Relaxation techniques for pain management in labour. *Cochrane Database of Systematic Reviews*, Issue 3. Art. No.: CD009514. DOI: 10.1002/14651858.CD009514.pub2.

Smith LA, Burns E, Cuthbert A. Parenteral opioids for maternal pain management in labour. *Cochrane Database of Systematic Reviews* 2018, Issue 6. Art. No.: CD007396. DOI: 10.1002/14651858.CD007396.pub3. Accessed 18 April 2021.

Spangler, A. K. et al. (2008) 'Belly models as teaching tools: What is their utility?', *Journal of Human Lactation*, 24(2), pp. 199–205. doi: 10.1177/0890334408316079.

Stark, M., Remynse, M. and Zwelling, E. (2016) 'Importance of the Birth Environment to Support Physiologic Birth.', *Journal of obstetric, gynecologic, and neonatal nursing: JOGNN*, 45(2), pp. 285–294. doi: 10.1016/j.jogn.2015.12.008.

Stephens TV, Payne M, Ball RO, Pencharz PB, Elango R. (2015). Protein requirements of healthy pregnant women during early and late gestation are higher than current recommendations. *The Journal of Nutrition*. 145(1):73-78.

Stillbirth Centre of Research Excellence (2021) 'Your baby's movements matter'. Safer Baby. Available at: <https://www.stillbirthcre.org.au/safer-baby-bundle/raising-awareness-and-improving-care-for-women-with-decreased-fetal-movements/>.

Stinson LF, Payne MS and Keelan JA (2018) A Critical Review of the Bacterial Baptism Hypothesis and the Impact of Cesarean Delivery on the Infant Microbiome. *Frontiers in Medicine*. 5 (135). Accessed online <https://www.ncbi-nlm-nih.gov.ezproxy.uws.edu.au/pmc/articles/PMC5945806/pdf/fmed-05-00135.pdf>

Suraci, N. et al. (2020) 'Improving labour progression among women with epidural anaesthesia following use of a birthing ball : a review of recent literature use of a birthing ball : a review of recent literature', *Journal Of Obstetrics and Gynaecology*, 40(4), pp. 491–494. doi: 10.1080/01443615.2019.1633519.

The Royal Women's Hospital (2018a) 'Abdominal muscle separation or diastasis'. Melbourne. Available at: <https://www.thewomens.org.au/images/uploads/fact-sheets/Abdominal-muscle-seperation-270719.pdf>.

The Royal Women's Hospital (2018b) *Induction of labour*. Melbourne. doi: 10.5694/j.1326-5377.1954.tb66970.x.

The Royal Women's Hospital, VIC. (n.d.). Managing pain in labour. Retrieved from <https://www.thewomens.org.au/health-information/pregnancy-and-birth/labour-birth/managing-pain-in-labour>

Tommys.org (2016) 'Reporting reduced fetal movements. What should I expect?' Available at: <https://www.tommys.org/pregnancy-information/pregnancy-symptom-checker/movements-matter-raising-awareness-fetal-movements>

Trotter S (2013). Why no baby skincare product should be advertised or promoted as 'suitable for newborn skin'. Midirs Midwifery Digest 23(2): p217-221.

<http://www.sharon trotter.org.uk/midirs2013.html>

Unicef UK (2021). Skin-to-skin contact. Accessed online:  
<https://www.unicef.org.uk/babyfriendly/baby-friendly-resources/implementing-standards-resources/skin-to-skin-contact/>

US Food and Drug Administration. (2020). Ultrasound Imaging. ONLINE  
<https://www.fda.gov/radiation-emitting-products/medical-imaging/ultrasound-imaging>

Vance, C. G., Dailey, D. L., Rakel, B. A., Sluka, K. A. (2014). Using TENS for pain control: the state of the evidence. Pain Management, 4(3), 197-209.  
[https://www.researchgate.net/publication/263322951\\_Using\\_TENS\\_for\\_pain\\_control\\_the\\_state\\_of\\_the\\_evidence](https://www.researchgate.net/publication/263322951_Using_TENS_for_pain_control_the_state_of_the_evidence)

van der Ploeg, J.M., Vervest, H.A.M., Liem, A.L., Schagen van Leeuwen, J.H. (1996). Transcutaneousnerve stimulation (TENS) during the first stage of labour: a randomized clinical trial. Pain, 68, 75-78. doi: [https://doi.org/10.1016/S0304-3959\(96\)03141-7](https://doi.org/10.1016/S0304-3959(96)03141-7)

Watts RW. Does pethidine still have a place in the management of labour pain?. Australian Prescriber 2004;27:34-5. <https://doi.org/10.18773/austprescr.2004.034>

Whitworth M, Bricker L, Mullan C. Ultrasound for fetal assessment in early pregnancy. Cochrane Database of Systematic Reviews 2015, Issue 7. Art. No.: CD007058. DOI: 10.1002/14651858.CD007058.pub3. Accessed 21 March 2021.

Woolcock, J. & Grivell, R. (2014). Noninvasive prenatal testing. *Australian Family Physician*. 43 (7), 432-434.

World Health Organisation (2014) *WHO recommendations for augmentation of labour*. Switzerland. Available at:

[https://www.who.int/reproductivehealth/publications/maternal\\_perinatal\\_health/augmentation-labour/en/](https://www.who.int/reproductivehealth/publications/maternal_perinatal_health/augmentation-labour/en/).

World Health Organisation. (2018). *WHO recommendations Intrapartum care for a positive childbirth experience*. Geneva: World Health Organization; 2018. Licence: CC BY-NC-SA 3.0 IGO. Accessed online

<https://apps.who.int/iris/bitstream/handle/10665/260178/9789241550215-eng.pdf;jsessionid=3577C95118F2E2DE8B62A1CCCD6F078B?sequence=1>